

# 1. Introduction

In 2011, total U.S. greenhouse gas emissions were 6,708.3 Tg or million metric tons CO<sub>2</sub> Eq. Total U.S. emissions have increased by 8.7 percent from 1990 to 2011, and emissions decreased from 2010 to 2011 by 1.5 percent (101.2 Tg CO<sub>2</sub> Eq.). The decrease from 2010 to 2011 was due to a decrease in the carbon intensity of fuels consumed to generate electricity due to a decrease in coal consumption, with increased natural gas consumption and a significant increase in hydropower used.

In 1992, the United States signed and ratified the United Nations Framework Convention on Climate Change (UNFCCC). As stated in Article 2 of the UNFCCC, “The ultimate objective of this Convention and any related legal instruments that the Conference of the Parties may adopt is to achieve, in accordance with the relevant provisions of the Convention, stabilization of greenhouse gas concentrations in the atmosphere at a level that would prevent dangerous anthropogenic interference with the climate system. Such a level should be achieved within a time-frame sufficient to allow ecosystems to adapt naturally to climate change, to ensure that food production is not threatened and to enable economic development to proceed in a sustainable manner.”<sup>20,21</sup>

Parties to the Convention, by ratifying, “shall develop, periodically update, publish and make available...national inventories of anthropogenic emissions by sources and removals by sinks of all greenhouse gases not controlled by the Montreal Protocol, using comparable methodologies...”<sup>22</sup> The United States views this report as an opportunity to fulfill these commitments under the UNFCCC.

In 1988, preceding the creation of the UNFCCC, the World Meteorological Organization (WMO) and the United Nations Environment Programme (UNEP) jointly established the Intergovernmental Panel on Climate Change (IPCC). The role of the IPCC is to assess on a comprehensive, objective, open and transparent basis the scientific, technical and socio-economic information relevant to understanding the scientific basis of risk of human-induced climate change, its potential impacts and options for adaptation and mitigation (IPCC 2003). Under Working Group 1 of the IPCC, nearly 140 scientists and national experts from more than thirty countries collaborated in the creation of the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC/UNEP/OECD/IEA 1997) to ensure that the emission inventories submitted to the UNFCCC are consistent and comparable between nations. The IPCC accepted the Revised 1996 IPCC Guidelines at its Twelfth Session (Mexico City, September 11-13, 1996). This report presents information in accordance with these guidelines. In addition, this Inventory is in accordance with the IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories and the Good Practice Guidance for Land Use, Land-Use Change, and Forestry, which further expanded upon the methodologies in the Revised 1996 IPCC Guidelines. The IPCC has also accepted the 2006 Guidelines for National Greenhouse Gas Inventories (IPCC 2006) at its Twenty-Fifth Session (Mauritius, April 2006). The 2006 IPCC Guidelines build on the previous bodies of work and includes new sources and gases “...as well as updates to the previously published methods whenever scientific and technical knowledge have improved since the previous guidelines were issued.” Many of the methodological improvements presented in the 2006 Guidelines have been adopted in this Inventory.

Overall, this inventory of anthropogenic greenhouse gas emissions provides a common and consistent mechanism through which Parties to the UNFCCC can estimate emissions and compare the relative contribution of individual sources, gases, and nations to climate change. The inventory provides a national estimate of sources and sinks for the United States, including all states and U.S. territories.<sup>23</sup> The structure of this report is consistent with the current UNFCCC Guidelines on Annual Inventories (UNFCCC 2006).

---

<sup>20</sup> The term “anthropogenic,” in this context, refers to greenhouse gas emissions and removals that are a direct result of human activities or are the result of natural processes that have been affected by human activities (IPCC/UNEP/OECD/IEA 1997).

<sup>21</sup> Article 2 of the Framework Convention on Climate Change published by the UNEP/WMO Information Unit on Climate Change. See <<http://unfccc.int>>. (UNEP/WMO 2000)

<sup>22</sup> Article 4(1)(a) of the United Nations Framework Convention on Climate Change (also identified in Article 12). Subsequent decisions by the Conference of the Parties elaborated the role of Annex I Parties in preparing national inventories. See <<http://unfccc.int>>.

<sup>23</sup> U.S. Territories include American Samoa, Guam, Puerto Rico, U.S. Virgin Islands, Wake Island, and other U.S. Pacific Islands.

[BEGIN BOX]

#### Box 1-1: Methodological approach for estimating and reporting U.S. emissions and sinks

In following the UNFCCC requirement under Article 4.1 to develop and submit national greenhouse gas emissions inventories, the emissions and sinks presented in this report are organized by source and sink categories and calculated using internationally-accepted methods provided by the IPCC.<sup>24</sup> Additionally, the calculated emissions and sinks in a given year for the U.S. are presented in a common manner in line with the UNFCCC reporting guidelines for the reporting of inventories under this international agreement.<sup>25</sup> The use of consistent methods to calculate emissions and sinks by all nations providing their inventories to the UNFCCC ensures that these reports are comparable. In this regard, U.S. emissions and sinks reported in this inventory report are comparable to emissions and sinks reported by other countries. Emissions and sinks provided in this inventory do not preclude alternative examinations, but rather this inventory report presents emissions and sinks in a common format consistent with how countries are to report inventories under the UNFCCC. The report itself follows this standardized format, and provides an explanation of the IPCC methods used to calculate emissions and sinks, and the manner in which those calculations are conducted.

On October 30, 2009, the U.S. Environmental Protection Agency (EPA) published a rule for the mandatory reporting of greenhouse gases (GHG) from large GHG emissions sources in the United States. Implementation of 40 CFR Part 98 is referred to as the Greenhouse Gas Reporting Program (GHGRP). 40 CFR part 98 applies to direct greenhouse gas emitters, fossil fuel suppliers, industrial gas suppliers, and facilities that inject CO<sub>2</sub> underground for sequestration or other reasons<sup>26</sup>. Reporting is at the facility level, except for certain suppliers of fossil fuels and industrial greenhouse gases. The GHGRP dataset and the data presented in this inventory report are complementary and, as indicated in the respective planned improvements sections in this report's chapters, EPA is analyzing the data for use, as applicable, to improve the national estimates presented in this inventory.

[END BOX]

## **1.1. Background Information**

### **Science**

For over the past 200 years, the burning of fossil fuels such as coal and oil, deforestation, and other sources have caused the concentrations of heat-trapping "greenhouse gases" to increase significantly in our atmosphere. These gases absorb some of the energy being radiated from the surface of the earth and trap it in the atmosphere, essentially acting like a blanket that makes the earth's surface warmer than it would be otherwise.

Greenhouse gases are necessary to life as we know it, because without them the planet's surface would be about 60 °F cooler than present. But, as the concentrations of these gases continue to increase in the atmosphere, the Earth's temperature is climbing above past levels. According to NOAA and NASA data, the Earth's average surface temperature has increased by about 1.2 to 1.4 °F since 1900. The ten warmest years on record (since 1850) have all occurred in the past 13 years (EPA 2009). Most of the warming in recent decades is very likely the result of human activities. Other aspects of the climate are also changing such as rainfall patterns, snow and ice cover, and sea level.

If greenhouse gases continue to increase, climate models predict that the average temperature at the Earth's surface could increase from 2.0 to 11.5 °F above 1990 levels by the end of this century (IPCC 2007). Scientists are certain that human activities are changing the composition of the atmosphere, and that increasing the concentration of

---

<sup>24</sup> See <<http://www.ipcc-nggip.iges.or.jp/public/index.html>>.

<sup>25</sup> See <[http://unfccc.int/national\\_reports/annex\\_i\\_ghg\\_inventories/national\\_inventories\\_submissions/items/5270.php](http://unfccc.int/national_reports/annex_i_ghg_inventories/national_inventories_submissions/items/5270.php)>

<sup>26</sup> See <<http://www.epa.gov/climatechange/emissions/ghgrulemaking.html>> and <<http://ghgdata.epa.gov/ghgp/main.do>>.

greenhouse gases will change the planet's climate. But they are not sure by how much it will change, at what rate it will change, or what the exact effects will be.<sup>27</sup>

### Greenhouse Gases

Although the Earth's atmosphere consists mainly of oxygen and nitrogen, neither plays a significant role in enhancing the greenhouse effect because both are essentially transparent to terrestrial radiation. The greenhouse effect is primarily a function of the concentration of water vapor, carbon dioxide (CO<sub>2</sub>), and other trace gases in the atmosphere that absorb the terrestrial radiation leaving the surface of the Earth (IPCC 2001). Changes in the atmospheric concentrations of these greenhouse gases can alter the balance of energy transfers between the atmosphere, space, land, and the oceans.<sup>28</sup> A gauge of these changes is called radiative forcing, which is a measure of the influence a factor has in altering the balance of incoming and outgoing energy in the Earth-atmosphere system (IPCC 2001). Holding everything else constant, increases in greenhouse gas concentrations in the atmosphere will produce positive radiative forcing (i.e., a net increase in the absorption of energy by the Earth).

*Climate change can be driven by changes in the atmospheric concentrations of a number of radiatively active gases and aerosols. We have clear evidence that human activities have affected concentrations, distributions and life cycles of these gases (IPCC 1996).*

Naturally occurring greenhouse gases include water vapor, CO<sub>2</sub>, methane (CH<sub>4</sub>), nitrous oxide (N<sub>2</sub>O), and ozone (O<sub>3</sub>). Several classes of halogenated substances that contain fluorine, chlorine, or bromine are also greenhouse gases, but they are, for the most part, solely a product of industrial activities. Chlorofluorocarbons (CFCs) and hydrochlorofluorocarbons (HCFCs) are halocarbons that contain chlorine, while halocarbons that contain bromine are referred to as bromofluorocarbons (i.e., halons). As stratospheric ozone depleting substances, CFCs, HCFCs, and halons are covered under the Montreal Protocol on Substances that Deplete the Ozone Layer. The UNFCCC defers to this earlier international treaty. Consequently, Parties to the UNFCCC are not required to include these gases in national greenhouse gas inventories.<sup>29</sup> Some other fluorine-containing halogenated substances—hydrofluorocarbons (HFCs), perfluorocarbons (PFCs), and sulfur hexafluoride (SF<sub>6</sub>)—do not deplete stratospheric ozone but are potent greenhouse gases. These latter substances are addressed by the UNFCCC and accounted for in national greenhouse gas inventories.

There are also several gases that, although they do not have a commonly agreed upon direct radiative forcing effect, do influence the global radiation budget. These tropospheric gases include carbon monoxide (CO), nitrogen dioxide (NO<sub>2</sub>), sulfur dioxide (SO<sub>2</sub>), and tropospheric (ground level) ozone O<sub>3</sub>. Tropospheric ozone is formed by two precursor pollutants, volatile organic compounds (VOCs) and nitrogen oxides (NO<sub>x</sub>) in the presence of ultraviolet light (sunlight). Aerosols are extremely small particles or liquid droplets that are often composed of sulfur compounds, carbonaceous combustion products, crustal materials and other human induced pollutants. They can affect the absorptive characteristics of the atmosphere. Comparatively, however, the level of scientific understanding of aerosols is still very low (IPCC 2001).

CO<sub>2</sub>, CH<sub>4</sub>, and N<sub>2</sub>O are continuously emitted to and removed from the atmosphere by natural processes on Earth. Anthropogenic activities, however, can cause additional quantities of these and other greenhouse gases to be emitted or sequestered, thereby changing their global average atmospheric concentrations. Natural activities such as respiration by plants or animals and seasonal cycles of plant growth and decay are examples of processes that only cycle carbon or nitrogen between the atmosphere and organic biomass. Such processes, except when directly or indirectly perturbed out of equilibrium by anthropogenic activities, generally do not alter average atmospheric greenhouse gas concentrations over decadal timeframes. Climatic changes resulting from anthropogenic activities, however, could have positive or negative feedback effects on these natural systems. Atmospheric concentrations of these gases, along with their rates of growth and atmospheric lifetimes, are presented in Table 1-1.

---

<sup>27</sup> For more information see <<http://www.epa.gov/climatechange/science>>

<sup>28</sup> For more on the science of climate change, see NRC (2001).

<sup>29</sup> Emissions estimates of CFCs, HCFCs, halons and other ozone-depleting substances are included in this document for informational purposes.

Table 1-1: Global Atmospheric Concentration, Rate of Concentration Change, and Atmospheric Lifetime (years) of Selected Greenhouse Gases

Atmospheric Variable	CO <sub>2</sub>	CH <sub>4</sub>	N <sub>2</sub> O	SF <sub>6</sub>	CF <sub>4</sub>
Pre-industrial atmospheric concentration	280 ppm	0.700 ppm	0.270 ppm	0 ppt	40 ppt
Atmospheric concentration	390 ppm	1.750-1.871 ppm <sup>a</sup>	0.322-0.323 ppm <sup>a</sup>	6.8-7.4 ppt	74 ppt
Rate of concentration change	1.4 ppm/yr	0.005 ppm/yr <sup>b</sup>	0.26%/yr	Linear <sup>c</sup>	Linear <sup>c</sup>
Atmospheric lifetime (years)	50-200 <sup>d</sup>	12 <sup>e</sup>	114 <sup>e</sup>	3,200	>50,000

Source: Pre-industrial atmospheric concentrations and rate of concentration changes for all gases are from IPCC (2007). The current atmospheric concentration for CO<sub>2</sub> is from NOAA/ESRL (2009).

<sup>a</sup> The range is the annual arithmetic averages from a mid-latitude Northern-Hemisphere site and a mid-latitude Southern-Hemisphere site for October 2006 through September 2007 (CDIAC 2009).

<sup>b</sup> The growth rate for atmospheric CH<sub>4</sub> has been decreasing from 1.4 ppb/yr in 1984 to less than 0 ppb/yr in 2001, 2004, and 2005.

<sup>c</sup> IPCC (2007) identifies the rate of concentration change for SF<sub>6</sub> and CF<sub>4</sub> as linear.

<sup>d</sup> No single lifetime can be defined for CO<sub>2</sub> because of the different rates of uptake by different removal processes.

<sup>e</sup> This lifetime has been defined as an “adjustment time” that takes into account the indirect effect of the gas on its own residence time.

A brief description of each greenhouse gas, its sources, and its role in the atmosphere is given below. The following section then explains the concept of GWPs, which are assigned to individual gases as a measure of their relative average global radiative forcing effect.

**Water Vapor (H<sub>2</sub>O).** Overall, the most abundant and dominant greenhouse gas in the atmosphere is water vapor. Water vapor is neither long-lived nor well mixed in the atmosphere, varying spatially from 0 to 2 percent (IPCC 1996). In addition, atmospheric water can exist in several physical states including gaseous, liquid, and solid. Human activities are not believed to affect directly the average global concentration of water vapor, but, the radiative forcing produced by the increased concentrations of other greenhouse gases may indirectly affect the hydrologic cycle. While a warmer atmosphere has an increased water holding capacity, increased concentrations of water vapor affects the formation of clouds, which can both absorb and reflect solar and terrestrial radiation. Aircraft contrails, which consist of water vapor and other aircraft emittants, are similar to clouds in their radiative forcing effects (IPCC 1999).

**Carbon Dioxide (CO<sub>2</sub>).** In nature, carbon is cycled between various atmospheric, oceanic, land biotic, marine biotic, and mineral reservoirs. The largest fluxes occur between the atmosphere and terrestrial biota, and between the atmosphere and surface water of the oceans. In the atmosphere, carbon predominantly exists in its oxidized form as CO<sub>2</sub>. Atmospheric CO<sub>2</sub> is part of this global carbon cycle, and therefore its fate is a complex function of geochemical and biological processes. CO<sub>2</sub> concentrations in the atmosphere increased from approximately 280 parts per million by volume (ppmv) in pre-industrial times to 389 ppmv in 2011, a 38.9 percent increase (IPCC 2007 and NOAA/ESRL 2012).<sup>30,31</sup> The IPCC definitively states that “the present atmospheric CO<sub>2</sub> increase is caused by anthropogenic emissions of CO<sub>2</sub>” (IPCC 2001). The predominant source of anthropogenic CO<sub>2</sub> emissions is the combustion of fossil fuels. Forest clearing, other biomass burning, and some non-energy production processes (e.g., cement production) also emit notable quantities of CO<sub>2</sub>. In its Fourth Assessment Report, the IPCC stated “most of the observed increase in global average temperatures since the mid-20<sup>th</sup> century is very likely due to the observed increased in anthropogenic greenhouse gas concentrations,” of which CO<sub>2</sub> is the most important (IPCC 2007).

**Methane (CH<sub>4</sub>).** CH<sub>4</sub> is primarily produced through anaerobic decomposition of organic matter in biological systems. Agricultural processes such as wetland rice cultivation, enteric fermentation in animals, and the decomposition of animal wastes emit CH<sub>4</sub>, as does the decomposition of municipal solid wastes. CH<sub>4</sub> is also emitted during the production and distribution of natural gas and petroleum, and is released as a by-product of coal mining and incomplete fossil fuel combustion. Atmospheric concentrations of CH<sub>4</sub> have increased by about 158

<sup>30</sup> The pre-industrial period is considered as the time preceding the year 1750 (IPCC 2001).

<sup>31</sup> Carbon dioxide concentrations during the last 1,000 years of the pre-industrial period (i.e., 750-1750), a time of relative climate stability, fluctuated by about ±10 ppmv around 280 ppmv (IPCC 2001).

percent since 1750, from a pre-industrial value of about 700 ppb to 1,750-1,871 ppb in 2010,<sup>32</sup> although the rate of increase has been declining. The IPCC has estimated that slightly more than half of the current CH<sub>4</sub> flux to the atmosphere is anthropogenic, from human activities such as agriculture, fossil fuel use, and waste disposal (IPCC 2007).

CH<sub>4</sub> is removed from the atmosphere through a reaction with the hydroxyl radical (OH) and is ultimately converted to CO<sub>2</sub>. Minor removal processes also include reaction with chlorine in the marine boundary layer, a soil sink, and stratospheric reactions. Increasing emissions of CH<sub>4</sub> reduce the concentration of OH, a feedback that may increase the atmospheric lifetime of CH<sub>4</sub> (IPCC 2001).

*Nitrous Oxide (N<sub>2</sub>O).* Anthropogenic sources of N<sub>2</sub>O emissions include agricultural soils, especially production of nitrogen-fixing crops and forages, the use of synthetic and manure fertilizers, and manure deposition by livestock; fossil fuel combustion, especially from mobile combustion; adipic (nylon) and nitric acid production; wastewater treatment and waste incineration; and biomass burning. The atmospheric concentration of N<sub>2</sub>O has increased by 19 percent since 1750, from a pre-industrial value of about 270 ppb to 322-323 ppb in 2010,<sup>33</sup> a concentration that has not been exceeded during the last thousand years. N<sub>2</sub>O is primarily removed from the atmosphere by the photolytic action of sunlight in the stratosphere (IPCC 2007).

*Ozone.* Ozone is present in both the upper stratosphere,<sup>34</sup> where it shields the Earth from harmful levels of ultraviolet radiation, and at lower concentrations in the troposphere,<sup>35</sup> where it is the main component of anthropogenic photochemical “smog.” During the last two decades, emissions of anthropogenic chlorine and bromine-containing halocarbons, such as CFCs, have depleted stratospheric ozone concentrations. This loss of ozone in the stratosphere has resulted in negative radiative forcing, representing an indirect effect of anthropogenic emissions of chlorine and bromine compounds (IPCC 1996). The depletion of stratospheric ozone and its radiative forcing was expected to reach a maximum in about 2000 before starting to recover. As of IPCC’s fourth assessment, “whether or not recently observed changes in ozone trends are already indicative of recovery of the global ozone layer is not yet clear” (IPCC 2007).

The past increase in tropospheric ozone, which is also a greenhouse gas, is estimated to provide the third largest increase in direct radiative forcing since the pre-industrial era, behind CO<sub>2</sub> and CH<sub>4</sub>. Tropospheric ozone is produced from complex chemical reactions of volatile organic compounds mixing with NO<sub>x</sub> in the presence of sunlight. The tropospheric concentrations of ozone and these other pollutants are short-lived and, therefore, spatially variable (IPCC 2001).

*Halocarbons, Perfluorocarbons, and Sulfur Hexafluoride.* Halocarbons are, for the most part, man-made chemicals that have both direct and indirect radiative forcing effects. Halocarbons that contain chlorine (CFCs, HCFCs, methyl chloroform, and carbon tetrachloride) and bromine (halons, methyl bromide, and hydrobromofluorocarbons [HFCs]) result in stratospheric ozone depletion and are therefore controlled under the Montreal Protocol on Substances that Deplete the Ozone Layer. Although CFCs and HCFCs include potent global warming gases, their net radiative forcing effect on the atmosphere is reduced because they cause stratospheric ozone depletion, which itself is an important greenhouse gas in addition to shielding the Earth from harmful levels of ultraviolet radiation. Under the Montreal Protocol, the United States phased out the production and importation of halons by 1994 and of CFCs by 1996. Under the Copenhagen Amendments to the Protocol, a cap was placed on the production and

---

<sup>32</sup> The range is the annual arithmetic averages from a mid-latitude Northern-Hemisphere site and a mid-latitude Southern-Hemisphere site for October 2006 through September 2007 (CDIAC 2010).

<sup>33</sup> The range is the annual arithmetic averages from a mid-latitude Northern-Hemisphere site and a mid-latitude Southern-Hemisphere site for October 2006 through September 2007 (CDIAC 2010).

<sup>34</sup> The stratosphere is the layer from the troposphere up to roughly 50 kilometers. In the lower regions the temperature is nearly constant but in the upper layer the temperature increases rapidly because of sunlight absorption by the ozone layer. The ozone-layer is the part of the stratosphere from 19 kilometers up to 48 kilometers where the concentration of ozone reaches up to 10 parts per million.

<sup>35</sup> The troposphere is the layer from the ground up to 11 kilometers near the poles and up to 16 kilometers in equatorial regions (i.e., the lowest layer of the atmosphere where people live). It contains roughly 80 percent of the mass of all gases in the atmosphere and is the site for most weather processes, including most of the water vapor and clouds.

importation of HCFCs by non-Article 5<sup>36</sup> countries beginning in 1996, and then followed by a complete phase-out by the year 2030. While ozone depleting gases covered under the Montreal Protocol and its Amendments are not covered by the UNFCCC, they are reported in this inventory under Annex 6.2 of this report for informational purposes.

HFCs, PFCs, and SF<sub>6</sub> are not ozone depleting substances, and therefore are not covered under the Montreal Protocol. They are, however, powerful greenhouse gases. HFCs are primarily used as replacements for ozone depleting substances but also emitted as a by-product of the HCFC-22 manufacturing process. Currently, they have a small aggregate radiative forcing impact, but it is anticipated that their contribution to overall radiative forcing will increase (IPCC 2001). PFCs and SF<sub>6</sub> are predominantly emitted from various industrial processes including aluminum smelting, semiconductor manufacturing, electric power transmission and distribution, and magnesium casting. Currently, the radiative forcing impact of PFCs and SF<sub>6</sub> is also small, but they have a significant growth rate, extremely long atmospheric lifetimes, and are strong absorbers of infrared radiation, and therefore have the potential to influence climate far into the future (IPCC 2001).

*Carbon Monoxide.* Carbon monoxide has an indirect radiative forcing effect by elevating concentrations of CH<sub>4</sub> and tropospheric ozone through chemical reactions with other atmospheric constituents (e.g., the hydroxyl radical, OH) that would otherwise assist in destroying CH<sub>4</sub> and tropospheric ozone. Carbon monoxide is created when carbon-containing fuels are burned incompletely. Through natural processes in the atmosphere, it is eventually oxidized to CO<sub>2</sub>. Carbon monoxide concentrations are both short-lived in the atmosphere and spatially variable.

*Nitrogen Oxides (NO<sub>x</sub>).* The primary climate change effects of nitrogen oxides (i.e., NO and NO<sub>2</sub>) are indirect and result from their role in promoting the formation of ozone in the troposphere and, to a lesser degree, lower stratosphere, where they have positive radiative forcing effects.<sup>37</sup> Additionally, NO<sub>x</sub> emissions from aircraft are also likely to decrease CH<sub>4</sub> concentrations, thus having a negative radiative forcing effect (IPCC 1999). Nitrogen oxides are created from lightning, soil microbial activity, biomass burning (both natural and anthropogenic fires) fuel combustion, and, in the stratosphere, from the photo-degradation of N<sub>2</sub>O. Concentrations of NO<sub>x</sub> are both relatively short-lived in the atmosphere and spatially variable.

*Nonmethane Volatile Organic Compounds (NMVOCs).* Non-CH<sub>4</sub> volatile organic compounds include substances such as propane, butane, and ethane. These compounds participate, along with NO<sub>x</sub>, in the formation of tropospheric ozone and other photochemical oxidants. NMVOCs are emitted primarily from transportation and industrial processes, as well as biomass burning and non-industrial consumption of organic solvents. Concentrations of NMVOCs tend to be both short-lived in the atmosphere and spatially variable.

*Aerosols.* Aerosols are extremely small particles or liquid droplets found in the atmosphere. They can be produced by natural events such as dust storms and volcanic activity, or by anthropogenic processes such as fuel combustion and biomass burning. Aerosols affect radiative forcing differently than greenhouse gases, and their radiative effects occur through direct and indirect mechanisms: directly by scattering and absorbing solar radiation; and indirectly by increasing droplet counts that modify the formation, precipitation efficiency, and radiative properties of clouds. Aerosols are removed from the atmosphere relatively rapidly by precipitation. Because aerosols generally have short atmospheric lifetimes, and have concentrations and compositions that vary regionally, spatially, and temporally, their contributions to radiative forcing are difficult to quantify (IPCC 2001).

The indirect radiative forcing from aerosols is typically divided into two effects. The first effect involves decreased droplet size and increased droplet concentration resulting from an increase in airborne aerosols. The second effect involves an increase in the water content and lifetime of clouds due to the effect of reduced droplet size on precipitation efficiency (IPCC 2001). Recent research has placed a greater focus on the second indirect radiative forcing effect of aerosols.

---

<sup>36</sup> Article 5 of the Montreal Protocol covers several groups of countries, especially developing countries, with low consumption rates of ozone depleting substances. Developing countries with per capita consumption of less than 0.3 kg of certain ozone depleting substances (weighted by their ozone depleting potential) receive financial assistance and a grace period of ten additional years in the phase-out of ozone depleting substances.

<sup>37</sup> NO<sub>x</sub> emissions injected higher in the stratosphere, primarily from fuel combustion emissions from high altitude supersonic aircraft, can lead to stratospheric ozone depletion.

Various categories of aerosols exist, including naturally produced aerosols such as soil dust, sea salt, biogenic aerosols, sulfates, and volcanic aerosols, and anthropogenically manufactured aerosols such as industrial dust and carbonaceous<sup>38</sup> aerosols (e.g., black carbon, organic carbon) from transportation, coal combustion, cement manufacturing, waste incineration, and biomass burning.

The net effect of aerosols on radiative forcing is believed to be negative (i.e., net cooling effect on the climate), although because they remain in the atmosphere for only days to weeks, their concentrations respond rapidly to changes in emissions.<sup>39</sup> Locally, the negative radiative forcing effects of aerosols can offset the positive forcing of greenhouse gases (IPCC 1996). “However, the aerosol effects do not cancel the global-scale effects of the much longer-lived greenhouse gases, and significant climate changes can still result” (IPCC 1996).

The IPCC’s Third Assessment Report notes that “the indirect radiative effect of aerosols is now understood to also encompass effects on ice and mixed-phase clouds, but the magnitude of any such indirect effect is not known, although it is likely to be positive” (IPCC 2001). Additionally, current research suggests that another constituent of aerosols, black carbon, has a positive radiative forcing, and that its presence “in the atmosphere above highly reflective surfaces such as snow and ice, or clouds, may cause a significant positive radiative forcing” (IPCC 2007). The primary anthropogenic emission sources of black carbon include diesel exhaust and open biomass burning.

## Global Warming Potentials

A global warming potential is a quantified measure of the globally averaged relative radiative forcing impacts of a particular greenhouse gas (see Table 1-2). It is defined as the ratio of the time-integrated radiative forcing from the instantaneous release of 1 kilogram (kg) of a trace substance relative to that of 1 kg of a reference gas (IPCC 2001). Direct radiative effects occur when the gas itself absorbs radiation. Indirect radiative forcing occurs when chemical transformations involving the original gas produce a gas or gases that are greenhouse gases, or when a gas influences other radiatively important processes such as the atmospheric lifetimes of other gases. The reference gas used is CO<sub>2</sub>, and therefore GWP-weighted emissions are measured in teragrams of CO<sub>2</sub> equivalent (Tg CO<sub>2</sub> Eq.).<sup>40</sup>

The relationship between gigagrams (Gg) of a gas and Tg CO<sub>2</sub> Eq. can be expressed as follows:

$$\text{Tg CO}_2 \text{ Eq} = (\text{Gg of gas}) \times (\text{GWP}) \times \left( \frac{\text{Tg}}{1,000 \text{ Gg}} \right)$$

where,

Tg CO<sub>2</sub> Eq. = Teragrams of CO<sub>2</sub> Equivalent

Gg = Gigagrams (equivalent to a thousand metric tons)

GWP = Global Warming Potential

Tg = Teragrams

GWP values allow for a comparison of the impacts of emissions and reductions of different gases. According to the IPCC, GWPs typically have an uncertainty of ±35 percent. The parties to the UNFCCC have also agreed to use GWPs based upon a 100-year time horizon, although other time horizon values are available.

*Greenhouse gas emissions and removals should be presented on a gas-by-gas basis in units of mass... In addition, consistent with decision 2/CP.3, Parties should report aggregate emissions and removals of greenhouse gases, expressed in CO<sub>2</sub> equivalent terms at summary inventory level, using GWP values provided by the IPCC in its Second Assessment Report... based on the effects of greenhouse gases over a*

<sup>38</sup> Carbonaceous aerosols are aerosols that are comprised mainly of organic substances and forms of black carbon (or soot) (IPCC 2001).

<sup>39</sup> Volcanic activity can inject significant quantities of aerosol producing sulfur dioxide and other sulfur compounds into the stratosphere, which can result in a longer negative forcing effect (i.e., a few years) (IPCC 1996).

<sup>40</sup> Carbon comprises 12/44<sup>ths</sup> of carbon dioxide by weight.

100-year time horizon.<sup>41</sup>

Greenhouse gases with relatively long atmospheric lifetimes (e.g., CO<sub>2</sub>, CH<sub>4</sub>, N<sub>2</sub>O, HFCs, PFCs, and SF<sub>6</sub>) tend to be evenly distributed throughout the atmosphere, and consequently global average concentrations can be determined. The short-lived gases such as water vapor, carbon monoxide, tropospheric ozone, ozone precursors (e.g., NO<sub>x</sub>, and NMVOCs), and tropospheric aerosols (e.g., SO<sub>2</sub> products and carbonaceous particles), however, vary regionally, and consequently it is difficult to quantify their global radiative forcing impacts. No GWP values are attributed to these gases that are short-lived and spatially inhomogeneous in the atmosphere.

Table 1-2: Global Warming Potentials and Atmospheric Lifetimes (Years) Used in this Report

Gas	Atmospheric Lifetime	GWP <sup>a</sup>
CO <sub>2</sub>	50-200	1
CH <sub>4</sub> <sup>b</sup>	12±3	21
N <sub>2</sub> O	120	310
HFC-23	264	11,700
HFC-32	5.6	650
HFC-125	32.6	2,800
HFC-134a	14.6	1,300
HFC-143a	48.3	3,800
HFC-152a	1.5	140
HFC-227ea	36.5	2,900
HFC-236fa	209	6,300
HFC-4310mee	17.1	1,300
CF <sub>4</sub>	50,000	6,500
C <sub>2</sub> F <sub>6</sub>	10,000	9,200
C <sub>4</sub> F <sub>10</sub>	2,600	7,000
C <sub>6</sub> F <sub>14</sub>	3,200	7,400
SF <sub>6</sub>	3,200	23,900

Source: (IPCC 1996)

<sup>a</sup> 100-year time horizon

<sup>b</sup> The GWP of CH<sub>4</sub> includes the direct effects and those indirect effects due to the production of tropospheric ozone and stratospheric water vapor. The indirect effect due to the production of CO<sub>2</sub> is not included.

[BEGIN BOX]

#### Box 1-2: The IPCC Fourth Assessment Report and Global Warming Potentials

In 2007, the IPCC published its Fourth Assessment Report (AR4), which provided an updated and more comprehensive scientific assessment of climate change. Within this report, the GWPs of several gases were revised relative to the SAR and the IPCC's Third Assessment Report (TAR) (IPCC 2001). Thus the GWPs used in this report have been updated twice by the IPCC; although the SAR GWPs are used throughout this report, it is interesting to review the changes to the GWPs and the impact such improved understanding has on the total GWP-weighted emissions of the United States. Since the SAR and TAR, the IPCC has applied an improved calculation of CO<sub>2</sub> radiative forcing and an improved CO<sub>2</sub> response function. The GWPs are drawn from IPCC/TEAP (2005) and the TAR, with updates for those cases where new laboratory or radiative transfer results have been published. Additionally, the atmospheric lifetimes of some gases have been recalculated. In addition, the values for radiative

<sup>41</sup> Framework Convention on Climate Change; <<http://unfccc.int/resource/docs/cop8/08.pdf>>; 1 November 2002; Report of the Conference of the Parties at its eighth session; held at New Delhi from 23 October to 1 November 2002; Addendum; Part One: Action taken by the Conference of the Parties at its eighth session; Decision -/CP.8; Communications from Parties Included in Annex I to the Convention: Guidelines for the Preparation of National Communications by Parties Included in Annex I to the Convention, Part 1: UNFCCC reporting guidelines on annual inventories; p. 7. (UNFCCC 2003)



forcing and lifetimes have been recalculated for a variety of halocarbons, which were not presented in the SAR. Table 1-3 presents the new GWPs, relative to those presented in the SAR.

Table 1-3: Comparison of 100-Year GWPs

Gas	SAR	TAR	AR4	Change from SAR	
				TAR	AR4
CO <sub>2</sub>	1	1	1	NC	0
CH <sub>4</sub> *	21	23	25	2	4
N <sub>2</sub> O	310	296	298	(14)	(12)
HFC-23	11,700	12,000	14,800	300	3,100
HFC-32	650	550	675	(100)	25
HFC-125	2,800	3,400	3,500	600	700
HFC-134a	1,300	1,300	1,430	NC	130
HFC-143a	3,800	4,300	4,470	500	670
HFC-152a	140	120	124	(20)	(16)
HFC-227ea	2,900	3,500	3,220	600	320
HFC-236fa	6,300	9,400	9,810	3,100	3,510
HFC-4310mee	1,300	1,500	1,640	200	340
CF <sub>4</sub>	6,500	5,700	7,390	(800)	890
C <sub>2</sub> F <sub>6</sub>	9,200	11,900	12,200	2,700	3,000
C <sub>4</sub> F <sub>10</sub>	7,000	8,600	8,860	1,600	1,860
C <sub>6</sub> F <sub>14</sub>	7,400	9,000	9,300	1,600	1,900
SF <sub>6</sub>	23,900	22,200	22,800	(1,700)	(1,100)

Source: (IPCC 2007, IPCC 2001)

NC (No Change)

Note: Parentheses indicate negative values.

\* The GWP of CH<sub>4</sub> includes the direct effects and those indirect effects due to the production of tropospheric ozone and stratospheric water vapor. The indirect effect due to the production of CO<sub>2</sub> is not included.

To comply with international reporting standards under the UNFCCC, official emission estimates are reported by the United States using SAR GWP values. The UNFCCC reporting guidelines for national inventories<sup>42</sup> were updated in 2002 but continue to require the use of GWPs from the SAR so that current estimates of aggregate greenhouse gas emissions for 1990 through 2011 are consistent and comparable with estimates developed prior to the publication of the TAR and AR4. For informational purposes, emission estimates that use the updated GWPs are presented in detail in Annex 6.1 of this report. All estimates provided throughout this report are also presented in unweighted units.

[END BOX]

## 1.2. Institutional Arrangements

The U.S. Environmental Protection Agency (EPA), in cooperation with other U.S. government agencies, prepares the Inventory of U.S. Greenhouse Gas Emissions and Sinks. A wide range of agencies and individuals are involved in supplying data to, reviewing, or preparing portions of the U.S. Inventory—including federal and state government authorities, research and academic institutions, industry associations, and private consultants.

Within EPA, the Office of Atmospheric Programs (OAP) is the lead office responsible for the emission calculations provided in the Inventory, as well as the completion of the National Inventory Report and the Common Reporting

<sup>42</sup> See <<http://unfccc.int/resource/docs/cop8/08.pdf>>.

Format tables. The Office of Transportation and Air Quality (OTAQ) is also involved in calculating emissions for the Inventory. While the U.S. Department of State officially submits the annual Inventory to the UNFCCC, EPA's OAP serves as the focal point for technical questions and comments on the U.S. Inventory. The staff of OAP and OTAQ coordinates the annual methodological choice, activity data collection, and emission calculations at the individual source category level. Within OAP, an inventory coordinator compiles the entire Inventory into the proper reporting format for submission to the UNFCCC, and is responsible for the collection and consistency of cross-cutting issues in the Inventory.

Several other government agencies contribute to the collection and analysis of the underlying activity data used in the Inventory calculations. Formal relationships exist between EPA and other U.S. agencies that provide official data for use in the Inventory. The U.S. Department of Energy's Energy Information Administration provides national fuel consumption data and the U.S. Department of Defense provides military fuel consumption and bunker fuels. Informal relationships also exist with other U.S. agencies to provide activity data for use in EPA's emission calculations. These include: the U.S. Department of Agriculture, the U.S. Geological Survey, the Federal Highway Administration, the Department of Transportation, the Bureau of Transportation Statistics, the Department of Commerce, the National Agricultural Statistics Service, and the Federal Aviation Administration. Academic and research centers also provide activity data and calculations to EPA, as well as individual companies participating in voluntary outreach efforts with EPA. Finally, the U.S. Department of State officially submits the Inventory to the UNFCCC each April.

### **1.3. Inventory Process**

EPA has a decentralized approach to preparing the annual U.S. Inventory, which consists of a National Inventory Report (NIR) and Common Reporting Format (CRF) tables. The Inventory coordinator at EPA is responsible for compiling all emission estimates and ensuring consistency and quality throughout the NIR and CRF tables. Emission calculations for individual sources are the responsibility of individual source leads, who are most familiar with each source category and the unique characteristics of its emissions profile. The individual source leads determine the most appropriate methodology and collect the best activity data to use in the emission calculations, based upon their expertise in the source category, as well as coordinating with researchers and contractors familiar with the sources. A multi-stage process for collecting information from the individual source leads and producing the Inventory is undertaken annually to compile all information and data.

### **Methodology Development, Data Collection, and Emissions and Sink Estimation**

Source leads at EPA collect input data and, as necessary, evaluate or develop the estimation methodology for the individual source categories. For most source categories, the methodology for the previous year is applied to the new "current" year of the Inventory, and inventory analysts collect any new data or update data that have changed from the previous year. If estimates for a new source category are being developed for the first time, or if the methodology is changing for an existing source category (e.g., the United States is implementing a higher Tiered approach for that source category), then the source category lead will develop a new methodology, gather the most appropriate activity data and emission factors (or in some cases direct emission measurements) for the entire time series, and conduct a special source-specific peer review process involving relevant experts from industry, government, and universities.

Once the methodology is in place and the data are collected, the individual source leads calculate emissions and sink estimates. The source leads then update or create the relevant text and accompanying annexes for the Inventory. Source leads are also responsible for completing the relevant sectoral background tables of the Common Reporting Format, conducting quality assurance and quality control (QA/QC) checks, and uncertainty analyses.

### **Summary Spreadsheet Compilation and Data Storage**

The inventory coordinator at EPA collects the source categories' descriptive text and Annexes, and also aggregates the emission estimates into a summary spreadsheet that links the individual source category spreadsheets together. This summary sheet contains all of the essential data in one central location, in formats commonly used in the Inventory document. In addition to the data from each source category, national trend and related data are also gathered in the summary sheet for use in the Executive Summary, Introduction, and Recent Trends sections of the Inventory report. Electronic copies of each year's summary spreadsheet, which contains all the emission and sink estimates for the United States, are kept on a central server at EPA under the jurisdiction of the Inventory

1 coordinator.

## 2 National Inventory Report Preparation

3 The NIR is compiled from the sections developed by each individual source lead. In addition, the inventory  
4 coordinator prepares a brief overview of each chapter that summarizes the emissions from all sources discussed in  
5 the chapters. The inventory coordinator then carries out a key category analysis for the Inventory, consistent with  
6 the IPCC Good Practice Guidance, IPCC Good Practice Guidance for Land Use, Land Use Change and Forestry,  
7 and in accordance with the reporting requirements of the UNFCCC. Also at this time, the Introduction, Executive  
8 Summary, and Recent Trends sections are drafted, to reflect the trends for the most recent year of the current  
9 Inventory. The analysis of trends necessitates gathering supplemental data, including weather and temperature  
10 conditions, economic activity and gross domestic product, population, atmospheric conditions, and the annual  
11 consumption of electricity, energy, and fossil fuels. Changes in these data are used to explain the trends observed in  
12 greenhouse gas emissions in the United States. Furthermore, specific factors that affect individual sectors are  
13 researched and discussed. Many of the factors that affect emissions are included in the Inventory document as  
14 separate analyses or side discussions in boxes within the text. Text boxes are also created to examine the data  
15 aggregated in different ways than in the remainder of the document, such as a focus on transportation activities or  
16 emissions from electricity generation. The document is prepared to match the specification of the UNFCCC  
17 reporting guidelines for National Inventory Reports.

## 18 Common Reporting Format Table Compilation

19 The CRF tables are compiled from individual tables completed by each individual source lead, which contain source  
20 emissions and activity data. The inventory coordinator integrates the source data into the UNFCCC's "CRF  
21 Reporter" for the United States, assuring consistency across all sectoral tables. The summary reports for emissions,  
22 methods, and emission factors used, the overview tables for completeness and quality of estimates, the recalculation  
23 tables, the notation key completion tables, and the emission trends tables are then completed by the inventory  
24 coordinator. Internal automated quality checks on the CRF Reporter, as well as reviews by the source leads, are  
25 completed for the entire time series of CRF tables before submission.

## 26 QA/QC and Uncertainty

27 QA/QC and uncertainty analyses are supervised by the QA/QC and Uncertainty coordinators, who have general  
28 oversight over the implementation of the QA/QC plan and the overall uncertainty analysis for the Inventory (see  
29 sections on QA/QC and Uncertainty, below). These coordinators work closely with the source leads to ensure that a  
30 consistent QA/QC plan and uncertainty analysis is implemented across all inventory sources. The inventory QA/QC  
31 plan, detailed in a following section, is consistent with the quality assurance procedures outlined by EPA and IPCC.

## 32 Expert and Public Review Periods

33 During the Expert Review period, a first draft of the document is sent to a select list of technical experts outside of  
34 EPA. The purpose of the Expert Review is to encourage feedback on the methodological and data sources used in  
35 the current Inventory, especially for sources which have experienced any changes since the previous Inventory.

36 Once comments are received and addressed, a second draft of the document is released for public review by  
37 publishing a notice in the U.S. Federal Register and posting the document on the EPA Web site. The Public Review  
38 period allows for a 30 day comment period and is open to the entire U.S. public.

## 39 Final Submittal to UNFCCC and Document Printing

40 After the final revisions to incorporate any comments from the Expert Review and Public Review periods, EPA  
41 prepares the final National Inventory Report and the accompanying Common Reporting Format Reporter database.  
42 The U.S. Department of State sends the official submission of the U.S. Inventory to the UNFCCC. The document is  
43 then formatted for printing, posted online, printed by the U.S. Government Printing Office, and made available for  
44 the public.

## 1.4. Methodology and Data Sources

Emissions of greenhouse gases from various source and sink categories have been estimated using methodologies that are consistent with the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC/UNEP/OECD/IEA 1997). In addition, the United States references the additional guidance provided in the IPCC Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories (IPCC 2000), the IPCC Good Practice Guidance for Land Use, Land-Use Change, and Forestry (IPCC 2003), and the 2006 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC 2006). To the extent possible, the present report relies on published activity and emission factor data. Depending on the emission source category, activity data can include fuel consumption or deliveries, vehicle-miles traveled, raw material processed, etc. Emission factors are factors that relate quantities of emissions to an activity.

The IPCC methodologies provided in the Revised 1996 IPCC Guidelines represent baseline methodologies for a variety of source categories, and many of these methodologies continue to be improved and refined as new research and data become available. This report uses the IPCC methodologies when applicable, and supplements them with other available methodologies and data where possible. Choices made regarding the methodologies and data sources used are provided in conjunction with the discussion of each source category in the main body of the report. Complete documentation is provided in the annexes on the detailed methodologies and data sources utilized in the calculation of each source category.

[BEGIN BOX]

### Box 1-3: IPCC Reference Approach

The UNFCCC reporting guidelines require countries to complete a "top-down" reference approach for estimating CO<sub>2</sub> emissions from fossil fuel combustion in addition to their "bottom-up" sectoral methodology. This estimation method uses alternative methodologies and different data sources than those contained in that section of the Energy chapter. The reference approach estimates fossil fuel consumption by adjusting national aggregate fuel production data for imports, exports, and stock changes rather than relying on end-user consumption surveys (see Annex 4 of this report). The reference approach assumes that once carbon-based fuels are brought into a national economy, they are either saved in some way (e.g., stored in products, kept in fuel stocks, or left unoxidized in ash) or combusted, and therefore the carbon in them is oxidized and released into the atmosphere. Accounting for actual consumption of fuels at the sectoral or sub-national level is not required.

[END BOX]

## 1.5. Key Categories

The IPCC's Good Practice Guidance (IPCC 2000) defines a key category as a "[source or sink category] that is prioritized within the national inventory system because its estimate has a significant influence on a country's total inventory of direct greenhouse gases in terms of the absolute level of emissions, the trend in emissions, or both."<sup>43</sup> By definition, key categories include those sources that have the greatest contribution to the absolute level of national emissions. In addition, when an entire time series of emission estimates is prepared, a thorough investigation of key categories must also account for the influence of trends and uncertainties of individual source and sink categories. This analysis culls out source and sink categories that diverge from the overall trend in national emissions. Finally, a qualitative evaluation of key categories is performed to capture any categories that were not identified in any of the quantitative analyses.

A Tier 1 approach, as defined in the IPCC's Good Practice Guidance (IPCC 2000), was implemented to identify the key categories for the United States. This analysis was performed twice; one analysis included sources and sinks

---

<sup>43</sup> See Chapter 7 "Methodological Choice and Recalculation" in IPCC (2000). <<http://www.ipcc-nggip.iges.or.jp/public/gp/gpgaum.htm>>

from the Land Use, Land-Use Change, and Forestry (LULUCF) sector, the other analysis did not include the LULUCF categories. Following the Tier 1 approach, a Tier 2 approach, as defined in the IPCC's Good Practice Guidance (IPCC 2000), was then implemented to identify any additional key categories not already identified in the Tier 1 assessment. This analysis, which includes each source category's uncertainty assessments (or proxies) in its calculations, was also performed twice to include or exclude LULUCF categories.

In addition to conducting Tier 1 and 2 level and trend assessments, a qualitative assessment of the source categories, as described in the IPCC's Good Practice Guidance (IPCC 2000), was conducted to capture any key categories that were not identified by either quantitative method. One additional key category, international bunker fuels, was identified using this qualitative assessment. International bunker fuels are fuels consumed for aviation or marine international transport activities, and emissions from these fuels are reported separately from totals in accordance with IPCC guidelines. If these emissions were included in the totals, bunker fuels would qualify as a key category according to the Tier 1 approach. The amount of uncertainty associated with estimation of emissions from international bunker fuels also supports the qualification of this source category as key, because it would qualify bunker fuels as a key category according to the Tier 2 approach.

Table 1-4 presents the key categories for the United States (including and excluding LULUCF categories) using emissions and uncertainty data in this report, and ranked according to their sector and global warming potential-weighted emissions in 2011. The table also indicates the criteria used in identifying these categories (i.e., level, trend, Tier 1, Tier 2, and/or qualitative assessments). Annex 1 of this report provides additional information regarding the key categories in the United States and the methodologies used to identify them.

Table 1-4: Key Categories for the United States (1990-2011)

IPCC Source Categories	Gas									Qual <sup>a</sup>	2011 Emissions (Tg CO <sub>2</sub> Eq.)
		Tier 1				Tier 2					
		Level Without LULUCF	Trend Without LULUCF	Level With LULUCF	Trend With LULUCF	Level Without LULUCF	Trend Without LULUCF	Level With LULUCF	Trend With LULUCF		
Energy											
CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Electricity Generation	CO <sub>2</sub>	•	•	•	•	•	•	•	•		1,722.7
CO <sub>2</sub> Emissions from Mobile Combustion: Road	CO <sub>2</sub>	•	•	•	•	•	•	•	•		1,466.7
CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Industrial	CO <sub>2</sub>	•	•	•	•	•	•	•	•		410.0
CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Electricity Generation	CO <sub>2</sub>	•	•	•	•	•	•	•	•		408.7
CO <sub>2</sub> Emissions from Stationary Combustion - Oil – Industrial	CO <sub>2</sub>	•	•	•	•	•	•	•	•		266.6
CO <sub>2</sub> Emissions from Stationary Combustion - Gas – Residential	CO <sub>2</sub>	•		•		•		•			255.7
CO <sub>2</sub> Emissions from Stationary Combustion - Gas - Commercial	CO <sub>2</sub>	•	•	•	•	•		•			171.1
CO <sub>2</sub> Emissions from Mobile Combustion: Aviation	CO <sub>2</sub>	•	•	•	•	•	•	•	•		145.2
CO <sub>2</sub> Emissions from Non-Energy Use of Fuels	CO <sub>2</sub>	•		•		•		•			130.6
CO <sub>2</sub> Emissions from	CO <sub>2</sub>	•	•	•	•	•	•	•	•		90.0

Stationary Combustion - Coal - Industrial												
CO <sub>2</sub> Emissions from Mobile Combustion: Other	CO <sub>2</sub>	•		•								82.5
CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Residential	CO <sub>2</sub>	•	•	•	•		•					73.5
CO <sub>2</sub> Emissions from Mobile Combustion: Marine	CO <sub>2</sub>	•		•								47.7
CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Commercial	CO <sub>2</sub>	•	•	•	•							46.6
CO <sub>2</sub> Emissions from Stationary Combustion - Oil - U.S. Territories	CO <sub>2</sub>	•	•	•	•		•					44.7
CO <sub>2</sub> Emissions from Natural Gas Systems	CO <sub>2</sub>	•	•	•	•	•	•	•				32.3
CO <sub>2</sub> Emissions from Stationary Combustion - Oil - Electricity Generation	CO <sub>2</sub>	•	•	•	•		•			•		26.6
CO <sub>2</sub> Emissions from Stationary Combustion - Coal - Commercial	CO <sub>2</sub>		•		•							5.0
Fugitive Emissions from Natural Gas Systems	CH <sub>4</sub>	•	•	•	•	•	•	•	•	•		139.6
Fugitive Emissions from Coal Mining	CH <sub>4</sub>	•	•	•	•	•	•	•	•	•		63.2
Fugitive Emissions from Petroleum Systems	CH <sub>4</sub>	•	•	•	•	•	•	•	•	•		31.5
Non-CO <sub>2</sub> Emissions from Stationary Combustion - Residential	CH <sub>4</sub>					•	•	•	•			3.5
Non-CO <sub>2</sub> Emissions from Stationary Combustion - Electricity Generation	N <sub>2</sub> O		•		•	•	•	•	•	•		17.9
N <sub>2</sub> O Emissions from Mobile Combustion: Road	N <sub>2</sub> O	•	•	•	•	•	•	•	•	•		14.4
Non-CO <sub>2</sub> Emissions from Stationary Combustion - Industrial	N <sub>2</sub> O						•					2.7
International Bunker Fuels <sup>c</sup>	Several									•		1.9

#### Industrial Processes

CO <sub>2</sub> Emissions from Iron and Steel Production & Metallurgical Coke Production	CO <sub>2</sub>	•	•	•	•	•	•	•	•			64.3
CO <sub>2</sub> Emissions from Cement Production	CO <sub>2</sub>	•	•	•	•							31.6
CO <sub>2</sub> Emissions from Ammonia Production	CO <sub>2</sub>		•		•							8.8
N <sub>2</sub> O Emissions from Adipic Acid Production	N <sub>2</sub> O		•		•							10.6
Emissions from Substitutes for Ozone Depleting Substances	HiGWP	•	•	•	•	•	•	•	•			121.7
SF <sub>6</sub> Emissions from Electrical Transmission and Distribution	HiGWP		•		•		•		•			7.0
HFC-23 Emissions from HCFC-22 Production	HiGWP	•	•	•	•		•		•			6.9

PFC Emissions from Aluminum Production	HiGWP		•		•						2.9
SF <sub>6</sub> Emissions from Magnesium Production and Processing	HiGWP		•								1.4
<b>Agriculture</b>											
CH <sub>4</sub> Emissions from Enteric Fermentation	CH <sub>4</sub>	•	•	•	•	•		•			137.4
CH <sub>4</sub> Emissions from Manure Management	CH <sub>4</sub>	•	•	•	•	•	•			•	52.0
CH <sub>4</sub> Emissions from Rice Cultivation	CH <sub>4</sub>					•		•			6.6
Direct N <sub>2</sub> O Emissions from Agricultural Soil Management	N <sub>2</sub> O	•		•		•		•			208.4
Indirect N <sub>2</sub> O Emissions from Applied Nitrogen	N <sub>2</sub> O	•		•		•	•	•			58.1
<b>Waste</b>											
CH <sub>4</sub> Emissions from Landfills	CH <sub>4</sub>	•	•	•	•	•	•	•	•		103.0
<b>Land Use, Land Use Change, and Forestry</b>											
CO <sub>2</sub> Emissions from Grassland Remaining Grassland	CO <sub>2</sub>			•	•			•	•		(9.0)
CO <sub>2</sub> Emissions from Landfilled Yard Trimmings and Food Scraps	CO <sub>2</sub>				•			•	•		(13.0)
CO <sub>2</sub> Emissions from Cropland Remaining Cropland	CO <sub>2</sub>			•	•			•	•		(14.6)
CO <sub>2</sub> Emissions from Urban Trees	CO <sub>2</sub>			•	•			•	•		(68.8)
CO <sub>2</sub> Emissions from Changes in Forest Carbon Stocks	CO <sub>2</sub>			•	•			•	•		(833.5)
CH <sub>4</sub> Emissions from Forest Fires	CH <sub>4</sub>				•			•	•		14.2
N <sub>2</sub> O Emissions from Forest Fires	N <sub>2</sub> O				•			•	•		11.6
<b>Subtotal Without LULUCF</b>											<b>6,716.5</b>
<b>Total Emissions Without LULUCF</b>											<b>6,851.2</b>
<b>Percent of Total Without LULUCF</b>											<b>98.0%</b>
<b>Subtotal With LULUCF</b>											<b>5,698.4</b>
<b>Total Emissions With LULUCF</b>											<b>5,829.4</b>
<b>Percent of Total With LULUCF</b>											<b>97.8%</b>

<sup>a</sup>Qualitative criteria.

<sup>b</sup>Emissions from this source not included in totals.

Note: Parentheses indicate negative values (or sequestration).

## 1 1.6. Quality Assurance and Quality Control (QA/QC)

2 As part of efforts to achieve its stated goals for inventory quality, transparency, and credibility, the United States has  
3 developed a quality assurance and quality control plan designed to check, document and improve the quality of its  
4 inventory over time. QA/QC activities on the Inventory are undertaken within the framework of the U.S. QA/QC  
5 plan, Quality Assurance/Quality Control and Uncertainty Management Plan for the U.S. Greenhouse Gas Inventory:  
6 Procedures Manual for QA/QC and Uncertainty Analysis.

7 Key attributes of the QA/QC plan are summarized in Figure 1-1. These attributes include:

- specific detailed procedures and forms that serve to standardize the process of documenting and archiving information, as well as to guide the implementation of QA/QC and the analysis of the uncertainty of the inventory estimates;
- expert review as well as QC—for both the inventory estimates and the Inventory (which is the primary vehicle for disseminating the results of the inventory development process). In addition, the plan provides for public review of the Inventory;
- both Tier 1 (general) and Tier 2 (source-specific) quality controls and checks, as recommended by IPCC Good Practice Guidance;
- consideration of secondary data quality and source-specific quality checks (Tier 2 QC) in parallel and coordination with the uncertainty assessment; the development of protocols and templates provides for more structured communication and integration with the suppliers of secondary information;
- record-keeping provisions to track which procedures have been followed, and the results of the QA/QC and uncertainty analysis, and feedback mechanisms for corrective action based on the results of the investigations, thereby providing for continual data quality improvement and guided research efforts;
- implementation of QA/QC procedures throughout the whole inventory development process—from initial data collection, through preparation of the emission estimates, to publication of the Inventory;
- a schedule for multi-year implementation; and
- promotion of coordination and interaction within the EPA, across Federal agencies and departments, state government programs, and research institutions and consulting firms involved in supplying data or preparing estimates for the Inventory. The QA/QC plan itself is intended to be revised and reflect new information that becomes available as the program develops, methods are improved, or additional supporting documents become necessary.

In addition, based on the national QA/QC plan for the Inventory, source-specific QA/QC plans have been developed for a number of sources. These plans follow the procedures outlined in the national QA/QC plan, tailoring the procedures to the specific text and spreadsheets of the individual sources. For each greenhouse gas emissions source or sink included in this Inventory, a minimum of a Tier 1 QA/QC analysis has been undertaken. Where QA/QC activities for a particular source go beyond the minimum Tier 1 level, further explanation is provided within the respective source category text.

The quality control activities described in the U.S. QA/QC plan occur throughout the inventory process; QA/QC is not separate from, but is an integral part of, preparing the inventory. Quality control—in the form of both good practices (such as documentation procedures) and checks on whether good practices and procedures are being followed—is applied at every stage of inventory development and document preparation. In addition, quality assurance occurs at two stages—an expert review and a public review. While both phases can significantly contribute to inventory quality, the public review phase is also essential for promoting the openness of the inventory development process and the transparency of the inventory data and methods.

The QA/QC plan guides the process of ensuring inventory quality by describing data and methodology checks, developing processes governing peer review and public comments, and developing guidance on conducting an analysis of the uncertainty surrounding the emission estimates. The QA/QC procedures also include feedback loops and provide for corrective actions that are designed to improve the inventory estimates over time.

Figure 1-1: U.S. QA/QC Plan Summary

## **1.7. Uncertainty Analysis of Emission Estimates – TO BE UPDATED**

Uncertainty estimates are an essential element of a complete and transparent emissions inventory. Uncertainty information is not intended to dispute the validity of the inventory estimates, but to help prioritize efforts to improve the accuracy of future inventories and guide future decisions on methodological choice. While the U.S. Inventory calculates its emission estimates with the highest possible accuracy, uncertainties are associated to a varying degree



with the development of emission estimates for any inventory. Some of the current estimates, such as those for CO<sub>2</sub> emissions from energy-related activities, are considered to have minimal uncertainty associated with them. For some other categories of emissions, however, a lack of data or an incomplete understanding of how emissions are generated increases the uncertainty surrounding the estimates presented. Despite these uncertainties, the UNFCCC reporting guidelines follow the recommendation in the 1996 IPCC Guidelines (IPCC/UNEP/OECD/IEA 1997) and require that countries provide single point estimates for each gas and emission or removal source category. Within the discussion of each emission source, specific factors affecting the uncertainty associated with the estimates are discussed.

Additional research in the following areas could help reduce uncertainty in the U.S. Inventory:

- *Incorporating excluded emission sources.* Quantitative estimates for some of the sources and sinks of greenhouse gas emissions are not available at this time. In particular, emissions from some land-use activities and industrial processes are not included in the inventory either because data are incomplete or because methodologies do not exist for estimating emissions from these source categories. See Annex 5 of this report for a discussion of the sources of greenhouse gas emissions and sinks excluded from this report.
- *Improving the accuracy of emission factors.* Further research is needed in some cases to improve the accuracy of emission factors used to calculate emissions from a variety of sources. For example, the accuracy of current emission factors applied to CH<sub>4</sub> and N<sub>2</sub>O emissions from stationary and mobile combustion is highly uncertain.
- *Collecting detailed activity data.* Although methodologies exist for estimating emissions for some sources, problems arise in obtaining activity data at a level of detail in which aggregate emission factors can be applied. For example, the ability to estimate emissions of SF<sub>6</sub> from electrical transmission and distribution is limited due to a lack of activity data regarding national SF<sub>6</sub> consumption or average equipment leak rates.

The overall uncertainty estimate for the U.S. greenhouse gas emissions inventory was developed using the IPCC Tier 2 uncertainty estimation methodology. Estimates of quantitative uncertainty for the overall greenhouse gas emissions inventory are shown below, in Table 1-5.

The IPCC provides good practice guidance on two approaches—Tier 1 and Tier 2—to estimating uncertainty for individual source categories. Tier 2 uncertainty analysis, employing the Monte Carlo Stochastic Simulation technique, was applied wherever data and resources permitted; further explanation is provided within the respective source category text and in Annex 7. Consistent with the IPCC Good Practice Guidance (IPCC 2000), over a multi-year timeframe, the United States expects to continue to improve the uncertainty estimates presented in this report.

Table 1-5: Estimated Overall Inventory Quantitative Uncertainty (Tg CO<sub>2</sub> Eq. and Percent)

Gas	2010 Emission Estimate <sup>a</sup> (Tg CO <sub>2</sub> Eq.)	Uncertainty Range Relative to Emission Estimate <sup>b</sup>				Mean <sup>c</sup> (Tg CO <sub>2</sub> Eq.)	Standard Deviation <sup>c</sup> (Tg CO <sub>2</sub> Eq.)
		(Tg CO <sub>2</sub> Eq.)		(%)			
		Lower Bound <sup>d</sup>	Upper Bound <sup>d</sup>	Lower Bound	Upper Bound		
CO <sub>2</sub>	5,706.0	5,570	5,958	-2%	4%	5,763	101
CH <sub>4</sub> <sup>e</sup>	666.5	578	751	-13%	13%	658	43
N <sub>2</sub> O <sup>e</sup>	306.2	265	431	-14%	41%	339	43
PFC, HFC & SF <sub>6</sub> <sup>e</sup>	140.3	138	156	-1%	11%	147	4
<b>Total</b>	<b>6,819.1</b>	<b>6,682</b>	<b>7,137</b>	<b>-2%</b>	<b>5%</b>	<b>6,906</b>	<b>117</b>
<b>Net Emissions (Sources and Sinks)</b>	<b>5,744.4</b>	<b>5,575</b>	<b>6,094</b>	<b>-3%</b>	<b>6%</b>	<b>5,830</b>	<b>133</b>

Notes:

<sup>a</sup> Emission estimates reported in this table correspond to emissions from only those source categories for which quantitative uncertainty was performed this year. Thus the totals reported in this table exclude approximately 2.8 Tg CO<sub>2</sub> Eq. of emissions for which quantitative uncertainty was not assessed. Hence, these emission estimates do not match the final total U.S. greenhouse gas emission estimates presented in this Inventory.

<sup>b</sup> The lower and upper bounds for emission estimates correspond to a 95 percent confidence interval, with the lower bound

corresponding to 2.5<sup>th</sup> percentile and the upper bound corresponding to 97.5<sup>th</sup> percentile.

<sup>c</sup> Mean value indicates the arithmetic average of the simulated emission estimates; standard deviation indicates the extent of deviation of the simulated values from the mean.

<sup>d</sup> The lower and upper bound emission estimates for the sub-source categories do not sum to total emissions because the low and high estimates for total emissions were calculated separately through simulations.

<sup>e</sup> The overall uncertainty estimates did not take into account the uncertainty in the GWP values for CH<sub>4</sub>, N<sub>2</sub>O and high GWP gases used in the inventory emission calculations for 2010.

Emissions calculated for the U.S. Inventory reflect current best estimates; in some cases, however, estimates are based on approximate methodologies, assumptions, and incomplete data. As new information becomes available in the future, the United States will continue to improve and revise its emission estimates. See Annex 7 of this report for further details on the U.S. process for estimating uncertainty associated with the emission estimates and for a more detailed discussion of the limitations of the current analysis and plans for improvement. Annex 7 also includes details on the uncertainty analysis performed for selected source categories.

## 1.8. Completeness

This report, along with its accompanying CRF reporter, serves as a thorough assessment of the anthropogenic sources and sinks of greenhouse gas emissions for the United States for the time series 1990 through 2011. Although this report is intended to be comprehensive, certain sources have been identified yet excluded from the estimates presented for various reasons. Generally speaking, sources not accounted for in this inventory are excluded due to data limitations or a lack of thorough understanding of the emission process. The United States is continually working to improve upon the understanding of such sources and seeking to find the data required to estimate related emissions. As such improvements are implemented, new emission sources are quantified and included in the Inventory. For a complete list of sources not included, see Annex 5 of this report.

## 1.9. Organization of Report

In accordance with the Revised 1996 IPCC Guidelines for National Greenhouse Gas Inventories (IPCC/UNEP/OECD/IEA 1997), and the 2006 UNFCCC Guidelines on Reporting and Review (UNFCCC 2006), this Inventory of U.S. Greenhouse Gas Emissions and Sinks is segregated into six sector-specific chapters, listed below in Table 1-6. In addition, chapters on Trends in Greenhouse Gas Emissions and Other information to be considered as part of the U.S. Inventory submission are included.

Table 1-6: IPCC Sector Descriptions

Chapter/IPCC Sector	Activities Included
Energy	Emissions of all greenhouse gases resulting from stationary and mobile energy activities including fuel combustion and fugitive fuel emissions.
Industrial Processes	By-product or fugitive emissions of greenhouse gases from industrial processes not directly related to energy activities such as fossil fuel combustion.
Solvent and Other Product Use	Emissions, of primarily NMVOCs, resulting from the use of solvents and N <sub>2</sub> O from product uses.
Agriculture	Anthropogenic emissions from agricultural activities except fuel combustion, which is addressed under Energy.
Land Use, Land-Use Change, and Forestry	Emissions and removals of CO <sub>2</sub> , CH <sub>4</sub> , and N <sub>2</sub> O from forest management, other land-use activities, and land-use change.
Waste	Emissions from waste management activities.

Source: (IPCC/UNEP/OECD/IEA 1997)

Within each chapter, emissions are identified by the anthropogenic activity that is the source or sink of the greenhouse gas emissions being estimated (e.g., coal mining). Overall, the following organizational structure is consistently applied throughout this report:

## **Chapter/IPCC Sector:** Overview of emission trends for each IPCC defined sector

**Source category:** Description of source pathway and emission trends.

**Methodology:** Description of analytical methods employed to produce emission estimates and identification of data references, primarily for activity data and emission factors.

**Uncertainty:** A discussion and quantification of the uncertainty in emission estimates and a discussion of time-series consistency.

**QA/QC and Verification:** A discussion on steps taken to QA/QC and verify the emission estimates, where beyond the overall U.S. QA/QC plan, and any key findings.

**Recalculations:** A discussion of any data or methodological changes that necessitate a recalculation of previous years' emission estimates, and the impact of the recalculation on the emission estimates, if applicable.

**Planned Improvements:** A discussion on any source-specific planned improvements, if applicable.

Special attention is given to CO<sub>2</sub> from fossil fuel combustion relative to other sources because of its share of emissions and its dominant influence on emission trends. For example, each energy consuming end-use sector (i.e., residential, commercial, industrial, and transportation), as well as the electricity generation sector, is described individually. Additional information for certain source categories and other topics is also provided in several Annexes listed in Table 1-7.

1 Table 1-7: List of Annexes

---

	ANNEX 1 Key Category Analysis
	ANNEX 2 Methodology and Data for Estimating CO <sub>2</sub> Emissions from Fossil Fuel Combustion
2.1.	Methodology for Estimating Emissions of CO <sub>2</sub> from Fossil Fuel Combustion
2.2.	Methodology for Estimating the Carbon Content of Fossil Fuels
2.3.	Methodology for Estimating Carbon Emitted from Non-Energy Uses of Fossil Fuels
	ANNEX 3 Methodological Descriptions for Additional Source or Sink Categories
3.1.	Methodology for Estimating Emissions of CH <sub>4</sub> , N <sub>2</sub> O, and Indirect Greenhouse Gases from Stationary Combustion
3.2.	Methodology for Estimating Emissions of CH <sub>4</sub> , N <sub>2</sub> O, and Indirect Greenhouse Gases from Mobile Combustion and Methodology for and Supplemental Information on Transportation-Related Greenhouse Gas Emissions
3.3.	Methodology for Estimating CH <sub>4</sub> Emissions from Coal Mining
3.4.	Methodology for Estimating CH <sub>4</sub> Emissions from Natural Gas Systems
3.5.	Methodology for Estimating CH <sub>4</sub> and CO <sub>2</sub> Emissions from Petroleum Systems
3.6.	Methodology for Estimating CO <sub>2</sub> and N <sub>2</sub> O Emissions from Incineration of Waste
3.7.	Methodology for Estimating Emissions from International Bunker Fuels used by the U.S. Military
3.8.	Methodology for Estimating HFC and PFC Emissions from Substitution of Ozone Depleting Substances
3.9.	Methodology for Estimating CH <sub>4</sub> Emissions from Enteric Fermentation
3.10.	Methodology for Estimating CH <sub>4</sub> and N <sub>2</sub> O Emissions from Manure Management
3.11.	Methodology for Estimating N <sub>2</sub> O Emissions and Soil Organic C Stock Changes from Agricultural Soil Management (Cropland and Grassland)
3.12.	Methodology for Estimating Net Carbon Stock Changes in Forest Lands Remaining Forest Lands
3.13.	Methodology for Estimating CH <sub>4</sub> Emissions from Landfills
	ANNEX 4 IPCC Reference Approach for Estimating CO <sub>2</sub> Emissions from Fossil Fuel Combustion
	ANNEX 5 Assessment of the Sources and Sinks of Greenhouse Gas Emissions Not Included
	ANNEX 6 Additional Information
6.1.	Global Warming Potential Values
6.2.	Ozone Depleting Substance Emissions
6.3.	Sulfur Dioxide Emissions
6.4.	Complete List of Source Categories
6.5.	Constants, Units, and Conversions
6.6.	Abbreviations
6.7.	Chemical Formulas
	ANNEX 7 Uncertainty
7.1.	Overview
7.2.	Methodology and Results
7.3.	Planned Improvements
7.4.	Additional Information on Uncertainty Analyses by Source

2

3